

Theory of 2D metal-insulator transition I: Zero Magnetic field

Konstantin Kladko

*Sun Microsystems, MS- CUP02-302,
901 San Antonio Road, Palo Alto, 94303 CA United States
e-mail:Konstantin.Kladko@sun.com*

We propose a scaling theory of 2D metal insulator transition discovered by Kravchenko and coworkers. In this theory conductance/resistance duality is an exact relation. The exponent of the stretched exponential in $\sigma(T)$ is determined by the temperature dependence of the coherence length. The quantum-chaotic nature of the transition is manifested in the square form of the phenomenological scaling function.

The 2D metal insulator transition (MIT) discovered by Kravchenko and coworkers is subject of wide interest. Nevertheless, to our knowledge, no satisfactory theory of such a transition has been proposed. Moreover, the discovery of such a transition puts in a serious question (in our opinion disproves) the validity of previous theories in this field. Here we propose a phenomenological scaling theory, which accounts for the following properties of the MIT at zero magnetic field. All of these properties were confirmed in different experiments on different systems. In our view, these properties summarize all interesting facts, measured in transport so far.

1. Resistance and conductance have a dual symmetry with respect to the critical point $\sigma = \sigma_0$.
2. At the critical point conductance does not depend on temperature.
3. All $\sigma(T)$ curves scale to the same universal curve.
4. $\sigma(T)$ shows approximate stretched exponential behavior $\simeq \exp CT^x$. The value of the stretched exponential x is not universal, and is different in different experiments.

To formulate a scaling theory, in the spirit of Landau, we need to guess the simplest phenomenological scaling relation for the conductance of the sample. It is clear, that at sample lengths larger, than the coherence length, quantum interference effects do not exist. Therefore, at such distances quantum scaling stops and conductance satisfies classical conductance addition laws. Let us imagine a square sample of length L , which has conductance $\sigma(L)$. Now we need to find a scaling equation which relates $\sigma(L)$ to $\sigma(2L)$, such as $\sigma(2L) = F(\sigma(L))$. The function $F(\sigma)$ is the main ingredient of the phenomenological theory. This function should be postulated by the comparison to the experiment. There is no need to derive it from anything else, since the underlying strongly interacting electron system is inherently chaotic and self-organized. We could, therefore, postulate $F(\sigma)$ without any derivations. We will, though, give a qualitative explanation for the chosen form of $F(\sigma)$.

In Landauer formalism (Fig.1) we phenomenologically relate $\sigma(L)$ to the number of conducting scattering channels, which go from the left side of the sample to the right side. Now let us consider squares A and B in Fig. 1. The conductance $\sigma(2L)$ will be then proportional to the probability for two L -channels, one from A and one from B , to meet and to form a long $2L$ -channel penetrating the larger square. Assuming, that excited state wave functions are very complicated, and channels meet chaotically, we can conclude, that the number of these $2L$ -channels will be proportional to the squared density of L -channels, i.e. to the squared conductance $\sigma(L)$. These arguments, therefore, lead to the following scaling equation

$$\sigma(2L) = C\sigma^2(L) \quad (1)$$

The function $F(\sigma)$ is, therefore, a quadratic function. We reiterate, that the above derivation is only a plausible analogy, which does not make an attempt to fully explain the complex structure of chaotic many-particle wave functions. It merely illustrates, that the quadratic $F(\sigma)$ is a right choice for the phenomenological theory. At the critical conductance resistance should be constant. Therefore, $C = 1/\sigma_0$, and

$$\sigma(2L) = \sigma^2(L)/\sigma_0 \quad (2)$$

Now, to find $\sigma(T)$ we need to integrate this equation from the microscopic length l to the coherence length $L_{coh}(T)$. Assuming, that the conductance at the microscopic length scale l is σ_l , the solution is

$$\sigma(T) = \sigma_0(\sigma_l/\sigma_0)^{L_{coh}(T)/l} \quad (3)$$

Here σ_l is the conductance at the microscopic length scale l . σ_l depends on carrier density. From here on we will measure σ in units of σ_0 . Then the above formula reads

$$\sigma(T) = \sigma_l^{L_{coh}(T)/l} \quad (4)$$

The above formula has a duality property, namely, transformation $\sigma_l \rightarrow 1/\sigma_l$ sends conductance into resistance and vice versa. This is, in our view, the duality which was observed in the experiment. If we assume, that σ_l smoothly depends on density near the transition, then near the transition one can always find such densities n and n' , that they will be in a duality relation. We note, that the duality relation is exact and does not depend on the specific temperature dependence of the coherence length. Now, if the coherence length goes as a power law of temperature $L_{coh} = A/T^\alpha$, that one usually assumes, one obtains a stretched exponential dependence of $\sigma(T)$.

$$\sigma(T) = \sigma_l^{\frac{A}{T^\alpha}} = e^{\frac{A}{T^\alpha} \ln(\sigma_l)} \quad (5)$$

Let us note that this dependence is approximate. Moreover, different stretched exponents observed in different experiments should correspond to different behaviors of the coherence length as a function of temperature in different systems. This resolves the question, why duality is universal and the value of the stretched exponential is, probably not. Nonuniversal values of α found in different experiments could mean, that the coherence length is set by external to the transition factors (phonons etc.).

We also, note, that in order for conducting phase to appear, σ_l , measured in units of σ_0 should become larger, than 1. The reason, why conducting phase never shows in lower mobility samples, is probably because σ_l there never gets large enough.

To incorporate magnetic field, one needs to include a phenomenological magnetic-field-dependent term into the right-hand side of the scaling equation. This will be done elsewhere. We believe, that this can be done consistently. We also believe, that phenomenological approach is the only approach which is meaningful in the case of the 2D MIT transition, which has inherently collective and chaotic nature.

I thank Sergej Kravchenko, Yaroslav Bazaliy and Pavel Kornilovich for enlightening comments and suggestions. This work has been supported by the Otto Hahn Medal Fellowship of the Max Planck Society, Germany.




Fig. 1

